

Table 4.31
Age-Adjusted Lung Cancer ^{1/} Death Rates
Calendar Years 1996-1998

	Deaths		Rate ^{2/}	
	Unadjusted	Adjusted ^{3/}	Unadjusted	Adjusted ^{3/}
<i>U.S. All Races (1997)</i>	153,276		37.4	
<i>All IHS Areas</i>	920	1,120	27.5	33.5
Aberdeen	97	100	53.4	55.1
Alaska	108	119	51.5	56.8
Albuquerque	16	16	9.5	9.5
Bemidji	141	179	70.1	88.9
Billings	53	59	49.5	55.1
California	60	91	19.7	30.1
Nashville	34	40	18.7	22.3
Navajo	27	27	5.7	5.7
Oklahoma	219	304	23.8	33.4
Phoenix	43	45	16.4	17.2
Portland	115	133	38.0	44.0
Tucson	7	7	10.0	10.0

^{1/} Lung cancer death includes deaths due to cancers of the trachea, bronchus and lung, ICD-9 codes 162.0 to 162.9.

^{2/} Age-adjusted rate per 100,000 population. Rates based on a small number of deaths should be interpreted with caution.

^{3/} Adjusted to compensate for misreporting of American Indian/Alaska Native race on the death certificate.